



Available online at www.ajcr.sinaweb.net

Apadana Journal of Clinical Research

Original Article

Estimation of health effects attributed to NO₂ exposure from the use of AirQ model in Ahvaz

Mohammad Javad Mohammadi¹, Gholamreza Goudarzi^{1,2*}, Ali K. Neisi^{1,2}, Kyvan Ahmadi Angali^{1,2}, Hamid Saki⁴, Ali Akbar Babaei², Bahram Mohammadi³, Hamid Dehdari Rad¹, Mehrdad Borhani⁴, Mehrdad Sharifi⁴

1. Assistant Professor of Genetics, Department of Environmental Health Engineering, School of Public Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran
2. Assistant Professor of Health sciences, Environmental Technologies Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran
3. Assistant Professor of Biology, Department of Biology PayamNour University Tehran unit, Tehran, Iran
4. Msc of Health Deputy of Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

Received 11 December 2012; received in revised form 12 January 2013; accepted 10 April 2013

Abstract

Background & objectives: AirQ software proved to be a valid and reliable tool to estimate the potential short term effects of air pollution. In this study the AirQ2.2.3 model was used to evaluate adverse health effects caused by nitrogen dioxide (NO₂) exposure in Ahvaz during 2009.

Materials & Methods: The AirQ, developed by the World Health Organization (WHO) to predict health endpoints attributed to criteria pollutants and permits the examination of various scenarios in which emission rates of pollutants are varied. Nitrogen dioxide (NO₂) reacts with ammonia, moisture, and other compounds to form small particles. These small particles penetrate deeply into sensitive parts of the lungs and can cause or worsen respiratory diseases, such as emphysema and bronchitis, and can aggravate existing heart disease, leading to increased hospital admissions and premature death.

Results: In the present study approximately 3% of total cardiovascular mortality, acute myocardial infarction and hospital admission of chronic obstructive pulmonary disease happened when the NO₂ concentration was more than 20 µg/m³.

Conclusions: Low percentage of observed health endpoints was associated to low concentration of measured NO₂. The lower level of RR value might achieve if some control strategies for reducing NO₂ emission were used. Therefore the higher RR value can depict mismanagement in urban air quality.

Keywords: Health endpoint; NO₂ concentration; AirQ model; Relative risk; Ahvaz

1. Introduction

Anthropogenic air pollution has been and continues to be viewed as a serious problem. Its seriousness lies in the fact that high, potentially harmful pollutant levels are

produced in environments which can be harmful for human health. According to the WHO estimations 800000 people were died each year because of cardiovascular and respiratory diseases, which are attributed to air pollution, in all over the world.

* Corresponding author: Gholamreza Goudarzi, Department of Environmental Health Engineering, School of Public Health, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran. Tel: +989163087159. E-mail: rgoodarzy@gmail.com

Approximately 150,000 of these deaths occur in the south of Asia. The result of these studies about short and long term effects of air pollution were estimated in the form of hospital admissions excess cases, increasing in a number of consultation with physician, asthma attacks, cardiopulmonary disease, death and number of the years lost (YOLL)¹⁻⁸. Nitrogen dioxide is produced in the combustion process which takes place mostly in transportation, power stations, heating plants and industrial processes. NO₂ and NO summation are called NO_x. The major source of NO_x is road traffic. Although 50% of total NO_x emission is produced by road traffic, this percentage can be higher in megacities and metropolitans⁹⁻¹¹. NO₂ can hurt lower parts of the respiratory system and at high concentrations, which may be occurred rarely at unprecedented industrial events and can lead to serious lung casualties. There are chronic and acute health effects which may be appeared by low and high NO₂ concentration exposure, respectively. High concentration of NO₂ can result in airway inflammation and this is why even healthy individuals show some anomalies in their respiratory system function. People with chronic obstructive pulmonary disease (COPD) or asthma attacks are more vulnerable at low concentration of NO₂¹²⁻¹³. Epidemiological studies have shown an association between NO₂ concentrations and health endpoints. In a multi-center case-cross-over study a 9.6 µg/m³ increase in the NO₂ concentration was associated with hospitalization for cardiovascular disease. In this study, increasing in hospital admission for cardiovascular disease, cardiac failure, ischemic heart disease and myocardial infarction were attributed to increase in the NO₂ concentration levels¹⁴. Again in another case-cross-over study, there was an association between NO₂ levels and mortality in patients who suffered from asthma exacerbation. In this study which was conducted in Spain, a 22.9 µg/m³ increase in the NO₂ concentration (minimum 5.2 µg/m³) was associated with mortality¹⁵. Positive associations were found between increases of 32 µg/m³ at temperatures above 20°C and hospital admissions for both primary intracerebral hemorrhage and ischemic stroke¹⁶. Next study has shown associations between hospital admissions for cardiovascular diseases and NO₂ for a 32 µg/m³ incremental increase¹⁷. Elevated odd ratios due to hospital admission for an increment of 10.5 µg/m³ NO₂ were observed among susceptible groups in a crossover study¹⁸. Recently, several studies

have reported adverse health effects below the current exposure limits for NO₂ particularly in both children and the elderly individuals. In a study with long term exposure (1-year average), 1.17, 1.57 were calculated as RR values with an increase of 16 µg/m³ in NO₂ concentration for all causes of death and cardiopulmonary disease, respectively. Again for the same increase of NO₂ concentrations in a long term exposure study (5-year average) relative risks of 1.19 and 1.74 were calculated for all causes of mortality and cardiopulmonary disease.

Patients and Methods

A specialized model, the air quality health impact assessment tool (AirQ_{2.2.3}), was used to assess the potential impacts of NO₂ exposure on human health in Ahvaz city during 2009. So, the mortality and morbidity of nitrogen dioxide in Ahvaz city in 2009 were calculated by AirQ_{2.2.3} utilizing relative risk and baseline incidence from WHO data. NO₂ data was taken from Ahvaz Department of Environment (ADoE). These data were in volumetric base. Health effects were being related to the mass of pollutants inhaled and which is why the AirQ model was on gravimetric basis. So, there was a conflict between AirQ model and ADoE data. Conversion between volumetric and gravimetric units (correction of temperature and pressure), coding, processing (averaging) and filtering are implemented for solving such problem.

Correction for non-standard temperature and pressure

Raw air quality monitoring data were in a Microsoft Office Excel spreadsheet. Therefore, all processing mechanisms such as correction, coding, averaging and filtering were performed in this software. The temperature and pressure are unlikely to be standard, so we also need to be able to convert gravimetric units at STP to other temperatures and pressures. At STP, we have 1 m³ containing a certain mass of material. When the temperature and pressure change, the volume of the gas changes but it still contains the same mass of material. Hence we need only to find the new volume from the Ideal Gas Equation

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

Where P_1 , V_1 and T_1 are the initial pressure, volume and absolute temperature and P_2 , V_2 and T_2 are the final pressure, volume and absolute temperature.

In our case:

$P_1 = 1$ atmosphere

$V_1 = 1 \text{ m}^3$

$T_1 = 273.15 \text{ K}$

And we need to find V_2 .

Therefore, rearranging equation below,

$$V_2 = \frac{T_2}{P_2} \times \frac{P_1 V_1}{T_1} = \frac{T_2}{273.15 \times P_2}$$

2-2- coding

We wrote a code in a column after date's column in spreadsheet excel. The objective for doing this was to

facilitate a mechanism of averaging. In the first cell of code column cell the formula for coding was:

$B3 = \text{Left}(A3, 5)$

To know what this formula means it should be noted that excel computes from date of Microsoft starting up till the first time of monitoring. Microsoft was founded in 1900. A count of the number of days from the first of 1900 till 03/21/2007 and the sum will be almost exactly 39,162. It was apparent by clicking on format cell/number/custom. The term of mm/dd/yyyyhh: main type part was corresponded to 39,162 in sample part.

Table1. coding on basis of date

| 0 | A(Date) | B(Code) | C(NO ₂ , µg/m ³) |
|----|------------------|---------|---|
| 1 | 03/21/2007 00:00 | 39162 | 164.01 |
| 2 | 03/21/2007 01:00 | 39162 | 174.57 |
| 3 | 03/21/2007 02:00 | 39162 | 165.98 |
| 21 | 03/21/2007 20:00 | 39162 | 111.59 |
| 22 | 03/21/2007 21:00 | 39162 | 110.12 |
| 23 | 03/21/2007 22:00 | 39162 | 107.91 |
| 24 | 03/21/2007 23:00 | 39162 | 73.47 |
| 25 | 03/22/2007 00:00 | 39163 | 81.77 |
| 26 | 03/22/2007 01:00 | 39163 | 53.90 |

Averaging

The routine method for providing 24 hours average is performed by clicking on f(x) icon. The insert function is appears. The routine way takes more time when there is a huge number of data. To obtain the daily average of the pollutant at each station in the city writing: (IF (B26<>B27; AVERAGE (C3:C26)) command would be necessary. According to the code column, the figure of code will be changed after 24 hours so that there is a special code for each day. The formula can generalize to the next cells and columns in both directions (horizontally and vertically). A slight change in the command sentence of formula would be recommended if it will spreads horizontally. For example above conditional sentence must alter to: (IF (B26<>B27;

AVERAGE (D3:D26)). Comparison of these two conditional formulas implies that C changes to D for providing average in next column which belongs to the levels of NO₂ at the adjacent station. By dragging up and down dragging the conditional sentence will act to compute daily mean of nitrogen dioxide in all stations. Huge numbers of cells (23 of 24 cells) are filled by the word "false" which means the first part of a conditional sentence is not correct. Therefore, mean value is placed on 24 number cells which are repeated in every 24 cell intervals.

Primary filtering

Primary filtering was used for deleting "false" cells. The command: Data/Filter/Auto filter/Does not equal to

false, is implemented to achieve 365 mean values of NO₂ concentrations at each station (Table 1).

Secondary filtering

According to model database, it is suggested to specify the highest and the lowest stations before using the secondary filtering. The station was considered as the highest when the mean value of this station is greater than the mean parameter at other stations and vice versa for the lowest. Therefore the lowest station should be selected i.e. the station that gives the lowest yearly average in the city for nitrogen dioxide concentration. In the next step, the column of all stations in the field of nitrogen dioxide concentration was built to make an appropriate background for secondary filtering. Then the air quality data were made from filtering on the column of all stations. The aim of secondary filtering on the average column of all stations was to find concentration intervals of NO₂ in 2007 year.

Results

The primary standard of nitrogen dioxide according to national ambient air quality standard (NAAQS) is 100 µg/m³ (NAAQS for air pollutant, Federal Register January 19, 2010). World health organization (AQG WHO) has recommended 40 and 200 µg/m³ as annual and hourly means of NO₂ guidelines, respectively (<http://www.euro.who.int/Document/E87950.pdf>).

Table 1 shows that annual mean of NO₂ in Ahvaz was 27 µg/m³ in 2009 which is lower than AQG WHO and also much lower than NAAQS standards. In view of NO₂ concentrations, Havashenasy and Edarekol were the highest and the lowest stations during this year respectively. The yearly average, summer mean, winter mean and 98 percentile of NO₂ concentrations in these stations is presented in table 2.

Table 2. NO₂ concentrations in micrograms µg/m³ for use in Model

| Parameter | lowest stations (Edarekol) | highest stations (Havashenasy) | Ahvaz |
|---------------|----------------------------|--------------------------------|-------|
| Annual mean | 18 | 66 | 27 |
| Summer mean | 16 | - | 6 |
| Winter mean | 19 | 72 | 43 |
| 98 percentile | 50 | 369 | 115 |

Relative risk and the estimated Attributable Proportion percentage for acute myocardial infarction were estimated in table 3. According to the model's default, the baseline incidence (BI) of this health endpoint for NO₂

was 132 per 10⁵ so the number of excess cases were estimated 9 at centerline of relative risk (RR=1.0036 and AP=0.6977%).

Table 3. Estimated relative risk indicators, and the component attributable to NO₂ cases attributable to acute myocardial infarction

| Indicator | Estimate | RR (Medium) | Estimated AP (%) | Estimated number of excess cases (persons) |
|-----------|----------|-------------|------------------|--|
| Down | | 1.0015 | 0.2919 | 3.7 |
| Mean | | 1.0036 | 0.6977 | 8.9 |
| Up | | 1.0084 | 1.6129 | 20.7 |




Figure1. Relationship between cumulative of acute MI versus NO₂ concentration

Figure 1 shows the cumulative number of acute MI versus NO₂ concentrations. Cumulative cases of this health endpoint were estimated by the model which was 9 in 2009. 60% of this number has occurred in the

days with concentrations lower than 70 µg/m³. It should be noted that 82% of above number are corresponded to the days with concentrations below 110 µg/m³.

Table 4. Estimated relative risk indicators, and the component attributable to NO₂ cases attributable to death from heart disease – cardiovascular

| Indicator | Estimate | RR (Medium) | Estimated AP (%) | Estimated number of excess cases (persons) |
|-----------|----------|-------------|------------------|--|
| Down | 1 | | 0.000 | 0.000 |
| Mean | 1.002 | | 0.3888 | 18.7 |
| Up | 1.004 | | 0.7746 | 37.3 |

Relative risk and the estimated Attributable Proportion percentage were estimated in table 4 for acute cardiovascular mortality. As mentioned earlier BI was taken from the model's default which is 497 per 10⁵ for this health endpoint. Therefore, estimated number of excess cases were calculated 19 at centerline of relative risk (RR=1.002 and AP=0.3888%).

Cumulative cases of cardiovascular mortality attributed to NO₂ concentrations has illustrated in Figure 2 with three ranges of relative risk. 37 persons were estimated by model as the total cumulative number of cardiovascular deaths within one year of exposure. 60% of these cases have occurred in the days with NO₂ levels not exceeding 70 µg/m³.




Figure 2.relationship between cumulative of cardiovascular mortality versus NO₂ concentration

Relative risk and the estimated Attributable Proportion percentage were estimated in table 4 for acute hospital admission of COPD. BI of this health endpoint is 101.4per

10⁵, the estimated number of excess cases was 8 at centerline of relative risk (RR=1.0038 and AP=0.7361%)

Table 5. Estimated relative risk indicators, and the component attributable to NO₂ cases attributable to chronic obstructive pulmonary disease (COPD)

| Indicator | Estimate | RR (Medium) | Estimated AP (%) | Estimated number of excess cases (persons) |
|-----------|----------|-------------|------------------|--|
| Down | | 1.0004 | 0.078 | 0.8 |
| Mean | | 1.0038 | 0.7361 | 7.2 |
| Up | | 1.0094 | 1.0094 | 17.7 |

Hospital admission of chronic obstructive pulmonary disease versus nitrogen dioxide concentration has been shown in figure 3. In this case the lowest line (RR=1.0004) is not attached to X axis and it shows a

subtle difference with figure 2. Estimated cases which attributed to NO₂ for hospital admission of COPD at lower, central and higher of RR were 1, 8 and 18, respectively (Table 5).




Figure 3- relationship between cumulative of hospital admission for COPD versus NO₂ concentration

Discussion

Figures 1 to 3 has illustrated NO₂ concentrations versus related health endpoint. As it has shown three ranges of relative risk based on model's default were considered in assessing health effects of NO₂. Furthermore BI values were also taken from default of the model. For our population of one million people and based on BI of 497 per 10⁵ people per year, some 4970 cardiovascular mortality cases can be expected annually, out of this number, 19 can be attributed to the NO₂ concentration above 20µg/m³. 82% of MI cases occurred in the days with pollutant not exceeding 110µg/m³. In cardiovascular mortality 77% of cases occurred in days with pollutant not exceeding 100 µg/m³ and 66% of COPD cases occurred in the days with pollutant not exceeding 80µg/m³. Although cardiovascular mortality had the lowest RR in centerline in comparison with MI and COPD, high number of this health endpoint cases was due to its higher BI value (497 per 10⁵). In the other hand, the greater number of acute cardiovascular mortality cases in comparison to MI cases was related to more BI for acute cardiovascular mortality. Even in comparison between MI and COPD, BI had a paramount of importance than RR. In the interpretation of RR it should be noted that 1 for this epidemiological parameter implies no impact on human health. In addition, severe effects can be expected when it exceeds from 1. Lack of cardiovascular mortality was referred to calculated RR value (1) and that is why the lowest line is attached to the X axis. It was also associated with being zero for both estimated AP and

estimated number of excess cases parameters. The lower level of RR value might be achieved if some control strategies for reducing NO₂ emission be used. Therefore the higher RR value can depict mismanagement in urban air quality. In the present study approximately 3 percent of total cardiovascular mortality, acute myocardial infarction and hospital admission of COPD happened when the NO₂ concentration was over 20µg/m³. Low percentage of observed health endpoints was associated to low concentration of measured NO₂ and as it mentioned NO₂ concentrations was lower than WHO and NAAQS guidelines.

Conclusion

In surgeries done on the nose, especially in aesthetic surgeries, efforts should be made to maintain the integrated structure of cartilage and as much as possible avoid manipulation and unnecessary cuts on the cartilage. In Suture technique, due to its non-destructiveness and minimum cartilage manipulation better results were obtained in the long term. It is recommended to apply VDD as the initial surgery method in sever deformities which cannot be corrected using Suture technique.

Acknowledgment

This paper is issued from M.SC thesis of Mohammad Javad Mohammadi and financial support

was provided by Ahvaz Jundishapur University of Medical Sciences.

References

- Barnett AG, Williams GM, Schwartz J, Best TL, Neller AH, Petroschevsky AL, et al. The effects of air pollution on hospitalizations for cardiovascular disease in elderly people in Australian and New Zealand cities. *Environ Health Perspect* 2006; 114(7): 1018-23.
- Bateson TF, Schwartz J. Who is sensitive to the effects of particulate air pollution on mortality? A case-cross over analysis of effect modifiers. *Epidemiology* 2004; 15(2): 143-9.
- Moll S, Gee D. Making sustainability accountable: eco-efficiency, resource productivity and innovation: proceedings of a workshop on the occasion of the Fifth Anniversary of the European Environment Agency (EEA) .Office for Official Publications of the European Communities; 1999.
- Oberdorster Gn, Finkelstein J, Ferin J, Gelein R, Johnston C, Godleski J, et al. Ultrafine particles as a potential environmental health hazard studies with model particles. *CHEST J* 1996; 109(3): 68-9.
- Gehring U, Heinrich J, Kramer U, Grote V, Hochadel M, Sugiri D, et al. Long-term exposure to ambient air pollution and cardiopulmonary mortality in women. *Epidemiology* 2006; 17(5): 545-51.
- Izzotti A, Parodi S, Quaglia A, Fare C, Vercelli M. The relationship between urban airborne pollution and short-term mortality: quantitative and qualitative aspects. *Eur J Epidemiol* 2000; 16(11): 1027-34.
- Berge E, Bartnicki J, Olendrzynski K, Tsyro S. Long-term trends in emissions and transboundary transport of acidifying air pollution in Europe. *J Environ Manag* 1999; 57(1): 31-50.
- Krzyzanowski M, Cohen A, Anderson R. Quantification of health effects of exposure to air pollution. *Ann Occup Environ Med* 2002; 59(12): 791-3.
- Pan Z, Molhave L, Kjaergaard SK. Effects on eyes and nose in humans after experimental exposure to airborne office dust. *Indoor air* 2000; 10(4): 237-45.
- Pope III CA, Burnett RT, Thun MJ, Calle EE, Krewski D, Ito K, et al. Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *Jama* 2002; 287(9): 1132-41.
- Reed M, Gigliotti A, McDonald J, Seagrave J, Seilkop S, Mauderly J. Health effects of subchronic exposure to environmental levels of diesel exhaust. *Inhal Toxicol* 2004; 16(4): 177-93.
- Salway A, Eggleston H, Goodwin J, Murrells T. UK emissions of air pollutants 1997.
- Sunyer J, Basagana X, Belmonte J, Anto J. Effect of nitrogen dioxide and ozone on the risk of dying in patients with severe asthma. *Thorax* 2002; 57(8): 687-93.
- Tsai SS, Goggins WB, Chiu HF, Yang CY. Evidence for an association between air pollution and daily stroke admissions in Kaohsiung, Taiwan. *Stroke* 2003; 34(11): 2612-6.
- Fielder R. Risk assessment of butadiene in ambient air; the approach used in the UK. *Toxicology* 1996; 113(1): 221-5.
- Van Leeuwen F. Update and revision of the WHO Air Quality Guidelines for Europe. *European EpiMarker* 1997; (1): 10-2.
- Krzyzanowski M. WHO air quality guidelines for Europe. *Environ Health Toxicol* 2008; 71(1): 47-50.
- Update and revision of the air quality guidelines for Europe: Meeting of the Working Group on Classical Air Pollutants. Copenhagen, WHO Regional Office for Europe 1995.